

nag_1d_cheb_interp_fit (e02afc)

1. Purpose

nag_1d_cheb_interp_fit (e02afc) computes the coefficients of a polynomial, in its Chebyshev series form, which interpolates (passes exactly through) data at a special set of points. Least-squares polynomial approximations can also be obtained.

2. Specification

```
#include <nag.h>
#include <nage02.h>

void nag_1d_cheb_interp_fit(Integer nplus1, double f[], double a[], NagError *fail)
```

3. Description

This routine computes the coefficients a_j , for $j = 1, 2, \dots, n + 1$, in the Chebyshev series

$$\frac{1}{2}a_1T_0(\bar{x}) + a_2T_1(\bar{x}) + a_3T_2(\bar{x}) + \dots + a_{n+1}T_n(\bar{x}),$$

which interpolates the data f_r at the points

$$\bar{x}_r = \cos((r-1)\pi/n), \quad r = 1, 2, \dots, n+1.$$

Here $T_j(\bar{x})$ denotes the Chebyshev polynomial of the first kind of degree j with argument \bar{x} . The use of these points minimizes the risk of unwanted fluctuations in the polynomial and is recommended when the data abscissae can be chosen by the user, e.g. when the data is given as a graph. For further advantages of this choice of points, see Clenshaw (1962).

In terms of the user's original variables, x say, the values of x at which the data f_r are to be provided are

$$x_r = \frac{1}{2}(x_{\max} - x_{\min}) \cos((r-1)\pi/n) + \frac{1}{2}(x_{\max} + x_{\min}), \quad r = 1, 2, \dots, n+1$$

where x_{\max} and x_{\min} are respectively the upper and lower ends of the range of x over which the user wishes to interpolate.

Truncation of the resulting series after the term involving a_{i+1} , say, yields a least-squares approximation to the data. This approximation, $p(\bar{x})$, say, is the polynomial of degree i which minimizes

$$\frac{1}{2}\epsilon_1^2 + \epsilon_2^2 + \epsilon_3^2 + \dots + \epsilon_n^2 + \frac{1}{2}\epsilon_{n+1}^2,$$

where the residual $\epsilon_r = p(\bar{x}_r) - f_r$, for $r = 1, 2, \dots, n+1$.

The method employed is based on the application of the three-term recurrence relation due to Clenshaw (1955) for the evaluation of the defining expression for the Chebyshev coefficients (see, for example, Clenshaw (1962)). The modifications to this recurrence relation suggested by Reinsch and Gentleman (see Gentleman (1969)) are used to give greater numerical stability.

For further details of the algorithm and its use see Cox (1974), Cox and Hayes (1973).

Subsequent evaluation of the computed polynomial, perhaps truncated after an appropriate number of terms, should be carried out using **nag_1d_cheb_eval (e02aec)**.

4. Parameters

nplus1

Input: the number $n + 1$ of data points (one greater than the degree n of the interpolating polynomial).

Constraint: **nplus1** ≥ 2 .

f[nplus1]

Input: for $r = 1, 2, \dots, n + 1$, **f**[$r - 1$] must contain f_r the value of the dependent variable (ordinate) corresponding to the value

$$\bar{x}_r = \cos\left(\frac{\pi(r-1)}{n}\right)$$

of the independent variable (abscissa) \bar{x} , or equivalently to the value

$$x_r = \frac{1}{2}(x_{\max} - x_{\min}) \cos(\pi(r-1)/n) + \frac{1}{2}(x_{\max} + x_{\min})$$

of the user's original variable x . Here x_{\max} and x_{\min} are respectively the upper and lower ends of the range over which the user wishes to interpolate.

a[nplus1]

Output: **a**[$j - 1$] is the coefficient a_j in the interpolating polynomial, for $j = 1, 2, \dots, n + 1$.

fail

The NAG error parameter, see the Essential Introduction to the NAG C Library.

5. Error Indications and Warnings

NE_INT_ARG_LT

On entry, **nplus1** must not be less than 2: **nplus1** = $\langle \text{value} \rangle$.

6. Further Comments

The time taken by the routine is approximately proportional to $(n + 1)^2 + 30$.

For choice of degree when using the routine for least-squares approximation, see the Chapter Introduction.

6.1. Accuracy

The rounding errors committed are such that the computed coefficients are exact for a slightly perturbed set of ordinates $f_r + \delta f_r$. The ratio of the sum of the absolute values of the δf_r to the sum of the absolute values of the f_r is less than a small multiple of $(n + 1)\epsilon$, where ϵ is the **machine precision**.

6.2. References

- Clenshaw C W (1955) A note on the summation of Chebyshev series *Math. Tables Aids Comput.* **9** 118–120.
- Clenshaw C W (1962) Mathematical tables *Chebyshev Series for Mathematical Functions* HMSO.
- Cox M G (1974) A data-fitting package for the non-specialist user *Software for Numerical Mathematics* (ed D J Evans) Academic Press.
- Cox M G and Hayes J G (1973) Curve fitting: a guide and suite of algorithms for the non-specialist user *NPL Report NAC 26* National Physical Laboratory.
- Gentleman W M (1969) An error analysis of Goertzel's (Watt's) method for computing Fourier coefficients *Comput. J.* **12** 160–165.

7. See Also

nag_1d_cheb_eval (e02aec)

8. Example

Determine the Chebyshev coefficients of the polynomial which interpolates the data \bar{x}_r, f_r , for $r = 1, 2, \dots, 11$, where $\bar{x}_r = \cos((r-1)\pi/10)$ and $f_r = e^{\bar{x}_r}$. Evaluate, for comparison with the values of f_r , the resulting Chebyshev series at \bar{x}_r , for $r = 1, 2, \dots, 11$.

The example program supplied is written in a general form that will enable polynomial interpolations of arbitrary data at the cosine points $\cos((r-1)\pi/n)$, for $r = 1, 2, \dots, n+1$ to be obtained for any n ($= \mathbf{nplus1}-1$). Note that `nag_1d_cheb_eval` (e02aec) is used to evaluate the interpolating polynomial. The program is self-starting in that any number of data sets can be supplied.

8.1. Program Text

```
/* nag_1d_cheb_interp_fit(e02afc) Example Program
 *
 * Copyright 1998 Numerical Algorithms Group.
 *
 * Mark 5, 1998.
 */

#include <nag.h>
#include <stdio.h>
#include <nag_stdlib.h>
#include <nage02.h>
#include <nagx01.h>
#include <math.h>

main()
{
#define NMAX 199
#define NP1MAX NMAX+1

    double d1;
    double xcap[NP1MAX];
    double f[NP1MAX];
    double pi, an[NP1MAX], fit;

    Integer i, j, n;
    Integer r;

    Vprintf("e02afc Example Program Results \n");

    /* Skip heading in data file */
    Vscanf("%*[\n]");

    pi = X01AAC;
    while((scanf("%ld",&n)) != EOF)
    {
        if (n > 0 && n <= NMAX)
        {
            pi = pi * 0.5 / (double) n;
            for (r = 0; r < n+1; ++r)
                Vscanf("%lf",&f[r]);

            for (r = 0; r < n+1; ++r)
            {
                i = r;
                /* The following method of evaluating xcap = cos(pi*i/n)
                 * ensures that the computed value has a small relative error
                 * and, moreover, is bounded in modulus by unity for all
                 * i = 0, 1, ..., n. (It is assumed that the sine routine
                 * produces a result with a small relative error for values
                 * of the argument between -PI/4 and PI/4).
                 */
                if (2*i <= n)
                {
                    d1 = sin(pi * i);
                    xcap[i] = 1.0 - d1 * d1 * 2.0;
                }
            }
        }
    }
}
```

```

    }
    else if (2*i > n * 3)
    {
        d1 = sin(piby2n * (n - i));
        xcap[i] = d1 * d1 * 2.0 - 1.0;
    }
    else
    {
        xcap[i] = sin(piby2n * (n - 2*i));
    }
}
e02afc(n+1, f, an, NAGERR_DEFAULT);
Vprintf("\n      Chebyshev \n");
Vprintf("    J    coefficient A(J) \n");
for (j = 0; j < n+1; ++j)
    Vprintf(" %3ld%14.7f\n", j+1, an[j]);
Vprintf("\n    R    Abscissa      Ordinate      Fit \n");
for (r = 0; r < n+1; ++r)
{
    e02aec(n+1, an, xcap[r], &fit, NAGERR_DEFAULT);
    Vprintf(" %3ld%11.4f%11.4f%11.4f\n", r+1, xcap[r], f[r], fit);
}
}
else
{
    Vprintf("Incorrect input value of n.\n");
    exit(EXIT_FAILURE);
}
}
exit(EXIT_SUCCESS);
}

```

8.2. Program Data

e02afc Example Program Data

```

10
2.7182
2.5884
2.2456
1.7999
1.3620
1.0000
0.7341
0.5555
0.4452
0.3863
0.3678

```

8.3. Program Results

e02afc Example Program Results

J	Chebyshev coefficient A(J)		
1	2.5320000		
2	1.1303095		
3	0.2714893		
4	0.0443462		
5	0.0055004		
6	0.0005400		
7	0.0000307		
8	-0.0000006		
9	-0.0000004		
10	0.0000049		
11	-0.0000200		

R	Abcissa	Ordinate	Fit
1	1.0000	2.7182	2.7182
2	0.9511	2.5884	2.5884
3	0.8090	2.2456	2.2456

4	0.5878	1.7999	1.7999
5	0.3090	1.3620	1.3620
6	0.0000	1.0000	1.0000
7	-0.3090	0.7341	0.7341
8	-0.5878	0.5555	0.5555
9	-0.8090	0.4452	0.4452
10	-0.9511	0.3863	0.3863
11	-1.0000	0.3678	0.3678
