

# NAG C Library Function Document

## nag\_1d\_cheb\_intg (e02ajc)

### 1 Purpose

nag\_1d\_cheb\_intg (e02ajc) determines the coefficients in the Chebyshev-series representation of the indefinite integral of a polynomial given in Chebyshev-series form.

### 2 Specification

```
void nag_1d_cheb_intg (Integer n, double xmin, double xmax, const double a[],
    Integer ia1, double qatm1, double aint[], Integer iaint1, NagError *fail)
```

### 3 Description

nag\_1d\_cheb\_intg (e02ajc) forms the polynomial which is the indefinite integral of a given polynomial. Both the original polynomial and its integral are represented in Chebyshev-series form. If supplied with the coefficients  $a_i$ , for  $i = 0, 1, \dots, n$ , of a polynomial  $p(x)$  of degree  $n$ , where

$$p(x) = \frac{1}{2}a_0 + a_1T_1(\bar{x}) + \dots + a_nT_n(\bar{x}),$$

the function returns the coefficients  $a'_i$ , for  $i = 0, 1, \dots, n+1$ , of the polynomial  $q(x)$  of degree  $n+1$ , where

$$q(x) = \frac{1}{2}a'_0 + a'_1T_1(\bar{x}) + \dots + a'_{n+1}T_{n+1}(\bar{x}),$$

and

$$q(x) = \int p(x) dx.$$

Here  $T_j(\bar{x})$  denotes the Chebyshev polynomial of the first kind of degree  $j$  with argument  $\bar{x}$ . It is assumed that the normalised variable  $\bar{x}$  in the interval  $[-1, +1]$  was obtained from the user's original variable  $x$  in the interval  $[x_{\min}, x_{\max}]$  by the linear transformation

$$\bar{x} = \frac{2x - (x_{\max} + x_{\min})}{x_{\max} - x_{\min}}$$

and that the user requires the integral to be with respect to the variable  $x$ . If the integral with respect to  $\bar{x}$  is required, set  $x_{\max} = 1$  and  $x_{\min} = -1$ .

Values of the integral can subsequently be computed, from the coefficients obtained, by using nag\_1d\_cheb\_eval2 (e02akc).

The method employed is that of Chebyshev-series (see Chapter 8 of Modern Computing Methods (1961)), modified for integrating with respect to  $x$ . Initially taking  $a_{n+1} = a_{n+2} = 0$ , the function forms successively

$$a'_i = \frac{a_{i-1} - a_{i+1}}{2i} \times \frac{x_{\max} - x_{\min}}{2}, \quad i = n+1, n, \dots, 1.$$

The constant coefficient  $a'_0$  is chosen so that  $q(x)$  is equal to a specified value, **qatm1**, at the lower end-point of the interval on which it is defined, i.e.,  $\bar{x} = -1$ , which corresponds to  $x = x_{\min}$ .

### 4 References

Modern Computing Methods (1961) Chebyshev-series *NPL Notes on Applied Science* **16** (2nd Edition) HMSO

## 5 Parameters

- 1: **n** – Integer *Input*  
*On entry:*  $n$ , the degree of the given polynomial  $p(x)$ .  
*Constraint:*  $n \geq 0$ .
- 2: **xmin** – double *Input*  
3: **xmax** – double *Input*  
*On entry:* the lower and upper end-points respectively of the interval  $[x_{\min}, x_{\max}]$ . The Chebyshev-series representation is in terms of the normalised variable  $\bar{x}$ , where
- $$\bar{x} = \frac{2x - (x_{\max} + x_{\min})}{x_{\max} - x_{\min}}.$$
- Constraint:* **xmax** > **xmin**.
- 4: **a[dim]** – const double *Input*  
**Note:** the dimension,  $dim$ , of the array **a** must be at least  $1 + n \times ia1$ .  
*On entry:* the Chebyshev coefficients of the polynomial  $p(x)$ . Specifically, element  $i \times ia1$  of **a** must contain the coefficient  $a_i$ , for  $i = 0, 1, \dots, n$ . Only these  $n + 1$  elements will be accessed.
- 5: **ia1** – Integer *Input*  
*On entry:* the index increment of **a**. Most frequently the Chebyshev coefficients are stored in adjacent elements of **a**, and **ia1** must be set to 1. However, if for example, they are stored in **a[0], a[3], a[6], ...**, then the value of **ia1** must be 3. See also Section 8.  
*Constraint:* **ia1**  $\geq 1$ .
- 6: **qatm1** – double *Input*  
*On entry:* the value that the integrated polynomial is required to have at the lower end-point of its interval of definition, i.e., at  $\bar{x} = -1$  which corresponds to  $x = x_{\min}$ . Thus, **qatm1** is a constant of integration and will normally be set to zero by the user.
- 7: **aint[dim]** – double *Output*  
**Note:** the dimension,  $dim$ , of the array **aint** must be at least  $1 + (n + 1) \times ia1$ .  
*On exit:* the Chebyshev coefficients of the integral  $q(x)$ . (The integration is with respect to the variable  $x$ , and the constant coefficient is chosen so that  $q(x_{\min})$  equals **qatm1**). Specifically, element  $i \times ia1$  of **aint** contains the coefficient  $a'_i$ , for  $i = 0, 1, \dots, n + 1$ .
- 8: **iaint1** – Integer *Input*  
*On entry:* the index increment of **aint**. Most frequently the Chebyshev coefficients are required in adjacent elements of **aint**, and **iaint1** must be set to 1. However, if, for example, they are to be stored in **aint[0], aint[3], aint[6], ...**, then the value of **iaint1** must be 3. See also Section 8.  
*Constraint:* **iaint1**  $\geq 1$ .
- 9: **fail** – NagError \* *Input/Output*  
The NAG error parameter (see the Essential Introduction).

## 6 Error Indicators and Warnings

### NE\_INT

*On entry,* **n** =  $\langle value \rangle$ .  
*Constraint:* **n**  $\geq 0$ .

On entry, **ia1** =  $\langle value \rangle$ .

Constraint: **ia1**  $\geq 1$ .

On entry, **iaint1** =  $\langle value \rangle$ .

Constraint: **iaint1**  $\geq 1$ .

## NE\_REAL\_2

On entry, **xmax**  $\leq$  **xmin**: **xmax** =  $\langle value \rangle$ , **xmin** =  $\langle value \rangle$ .

## NE\_BAD\_PARAM

On entry, parameter  $\langle value \rangle$  had an illegal value.

## NE\_INTERNAL\_ERROR

An internal error has occurred in this function. Check the function call and any array sizes. If the call is correct then please consult NAG for assistance.

## 7 Accuracy

In general there is a gain in precision in numerical integration, in this case associated with the division by  $2i$  in the formula quoted in Section 3.

## 8 Further Comments

The time taken is approximately proportional to  $n + 1$ .

The increments **ia1**, **iaint1** are included as parameters to give a degree of flexibility which, for example, allows a polynomial in two variables to be integrated with respect to either variable without rearranging the coefficients.

## 9 Example

Suppose a polynomial has been computed in Chebyshev-series form to fit data over the interval  $[-0.5, 2.5]$ . The following program evaluates the integral of the polynomial from 0.0 to 2.0. (For the purpose of this example, **xmin**, **xmax** and the Chebyshev coefficients are simply supplied. Normally a program would read in or generate data and compute the fitted polynomial).

### 9.1 Program Text

```
/* nag_ld_chheb_intg (e02ajc) Example Program.
 *
 * Copyright 2001 Numerical Algorithms Group.
 *
 * Mark 7, 2001.
 */

#include <stdio.h>
#include <nag.h>
#include <nag_stdlib.h>
#include <nage02.h>

int main(void)
{
    /* Initialized data */
    const double xmin = -0.5;
    const double xmax = 2.5;
    const double a[7] = { 2.53213, 1.13032, 0.2715, 0.04434, 0.00547, 5.4e-4, 4e-5 };

    /* Scalars */
    double ra, rb, result, xa, xb, zero;
    Integer exit_status, n, one;
    NagError fail;
```

```

/* Arrays */
double *aint = 0;

INIT_FAIL(fail);
exit_status = 0;
Vprintf("e02ajc Example Program Results\n");

n = 6;
zero = 0.0;
one = 1;

/* Allocate memory */
if ( !(aint = NAG_ALLOC(n + 2, double)) )
{
    Vprintf("Allocation failure\n");
    exit_status = -1;
    goto END;
}

e02ajc(n, xmin, xmax, a, one, zero, aint, one, &fail);
if (fail.code != NE_NOERROR)
{
    Vprintf("Error from e02ajc.\n%s\n", fail.message);
    exit_status = 1;
    goto END;
}

xa = 0.0;
xb = 2.0;
e02akc(n+1, xmin, xmax, aint, one, xa, &ra, &fail);
if (fail.code != NE_NOERROR)
{
    Vprintf("Error from e02akc.\n%s\n", fail.message);
    exit_status = 1;
    goto END;
}

e02akc(n+1, xmin, xmax, aint, one, xb, &rb, &fail);
if (fail.code != NE_NOERROR)
{
    Vprintf("Error from e02akc.\n%s\n", fail.message);
    exit_status = 1;
    goto END;
}

result = rb - ra;
Vprintf("\n");
Vprintf("Value of definite integral is %10.4f\n", result);
END:
if (aint) NAG_FREE(aint);

return exit_status;
}

```

## 9.2 Program Data

None.

## 9.3 Program Results

e02ajc Example Program Results

Value of definite integral is      2.1515

---